

Get them
IN ON TIME
 through the 1st year

Birth	
HepB	

Age 2 months	Not earlier than 6 weeks of age
RGE (Rotavirus)	
HepB	
DTaP	
Hib	Not earlier than 6 weeks of age
Polio (IPV)	
PCV (Pneumo)	

Age 4 months	Interval from previous dose
RGE (Rotavirus)	
DTaP	1-2 months
Hib	1-2 months
Polio (IPV)	1-2 months
PCV (Pneumo)	1-2 months

Age 6 months	Interval from previous dose
RGE (Rotavirus)	
HepB	at least 4 months after first dose
DTaP	1-2 months
Hib	1-2 months
Polio (IPV)	1-2 months
PCV (Pneumo)	1-2 months

Age 12 months	Interval from previous dose
DTaP	6 months after the third dose
Hib	2 months
PCV (Pneumo)	8 weeks and on or after first birthday
MMR	On or after first birthday
Varicella	On or after first birthday
HepA	2nd dose 6 months later



Every Fall: Flu Vaccine

- All children 6 months to 5 years of age.
- All children 6 months and older with asthma, diabetes, or other chronic high-risk condition.
- Others in the household of both the above.

Note: Children 6 months through 8 years old need 2 doses, one month apart, in their first year of flu shots.

Before Kindergarten:

- DTaP, Polio, MMR#2, also, unless already given, HepB and Varicella

Indiana's "IN ON TIME" Childhood Immunization Schedule is compatible with the recommendation of the Advisory Committee on Immunization Practices (ACIP) of the US Centers for Disease Control and Prevention (CDC), the American Academy of Pediatrics (AAP) and the American Academy of Family Physicians (AAFP). If you have any questions, call the Indiana State Department of Health, 800-701-0704.

Combined vaccines may decrease the number of shots.

Updated on March 2, 2007